

Periodic instanton bifurcations and thermal transition rate

A. N. Kuznetsov and P. G. Tinyakov

*Institute for Nuclear Research of the Russian Academy of Sciences,
60th October Anniversary prospect 7a, Moscow 117312, Russia.*

February 1, 2008

Abstract

It is shown that periodic instanton solutions may have bifurcations which qualitatively change the behaviour of the finite temperature transition rate. These bifurcations are studied numerically in a quantum mechanical model and in the massive two-dimensional sigma model with the Skyrme term. General rules which determine the behaviour of periodic instantons near bifurcations are derived and applied to the Electroweak Theory.

In weakly coupled theories, tunneling transitions can often be associated with classical solutions to Euclidean equations of motion. A special class of such solutions, periodic solutions with two turning points (periodic instantons), are believed to dominate the finite energy [1] and, under certain conditions, the finite temperature transition rate [2].

Periodic instantons naturally emerge in the context of tunneling from a micro-canonical ensemble with fixed energy E . The probability of tunneling from such initial state can be written as follows,

$$P(E) = \sum_{i,f} |\langle f | \hat{S} \hat{P}_E | i \rangle|^2, \quad (1)$$

where $\hat{S}$ is the S-matrix, $\hat{P}_E$ is projector onto subspace of fixed energy E , while the states $|i\rangle$ and $|f\rangle$ are excitations above two vacua lying on different sides of the barrier. Eq.(1) has a path integral representation [1]. In the semi-classical limit the path integral is expected to be saturated by a single solution to Euclidean field equations possessing two turning points at $t = 0$ and $t = i\tau/2$ and, hence, periodic in Euclidean time with period τ — the periodic instanton. With exponential accuracy the probability $P(E)$ is

$$P(E) \sim e^{W(E)} = e^{E\tau - S(\tau)}, \quad (2)$$

where $S(\tau)$ is the action of the periodic instanton per period and τ is related to the energy E in the standard way,

$$E = \frac{\partial S}{\partial \tau}. \quad (3)$$

Eq.(2) holds at $E < E_{sph}$, where E_{sph} is the height of the potential barrier¹. At $E > E_{sph}$ the transition probability is not exponentially suppressed, $P(E) \sim 1$.

The thermal rate is constructed from $P(E)$ by averaging with Boltzmann exponent. The transition rate for Gibbs ensemble at temperature $T = 1/\beta$ equals

$$\begin{aligned} \Gamma(\beta) &= \int_0^\infty dE e^{-\beta E} P(E) \\ &\sim \int_0^\infty dE e^{-\beta E + W(E)}. \end{aligned} \quad (4)$$

In the weak coupling limit, $g \ll 1$, and temperatures $T \ll 1/g$ the integral over E can be calculated by the steepest descent method. Formally, the saddle point condition in eq. (4) reads

$$\beta = \tau(E), \quad (5)$$

where $\tau(E)$ is the period of the periodic instanton. Thus, only periodic instantons with the period equal to inverse temperature can dominate the thermal rate. The relevance of a particular saddle point depends on the sign of $\partial^2 W / \partial E^2$. As we will see below, in many models this quantity may change sign at certain values of temperature. As a result, in different temperature ranges the transition rate is dominated by different solutions.

The exact analytic form of the periodic instanton is known only in one-dimensional quantum mechanics. In field theory models it can be found either approximately at low energies [1] or numerically [4, 5, 6]. However, the qualitative picture of periodic instanton behaviour can often be obtained from its low and high energy limits by making use of a few general rules it obeys. This picture in many cases turns out to be rather complicated. Namely, periodic instantons may have bifurcations which must be taken into account in the calculation of the finite temperature transition rate.

Consider first the high energy limit $E \sim E_{sph}$. At $E = E_{sph}$ the periodic instanton reduces to the sphaleron. Since the sphaleron is the static solution to the field equations, it is “periodic” with any period. The action per period of this solution is equal to $E_{sph}\tau$ while $W(E_{sph})$ is identically zero.

Close to the sphaleron, $E - E_{sph} \ll E_{sph}$, the periodic instanton can be represented as the sum of the sphaleron and an (almost linear) oscillation in its negative eigenmode. In the limit $E \rightarrow E_{sph}$ the period of the oscillation approaches $\tau_{sph} = 2\pi/\omega_-$, where ω_- is the sphaleron negative eigenvalue, while the amplitude goes to zero and the periodic instanton merges into the sphaleron. The behaviour of the period $\tau(E)$

¹In the Electroweak Theory the top of the barrier is associated with the sphaleron solution [3], which is the origin of the notation E_{sph} .




Figure 1: Two possible types of smooth behaviour of the periodic instanton action S as a function of its period τ . The straight line represents the sphaleron with the action $S = E_{sph}\tau$.

in the vicinity of the sphaleron is determined by non-linear effects. Depending on particular model the period τ can be larger or smaller than τ_{sph} ; correspondingly, the action $S(\tau)$ is smaller or larger than $E_{sph}\tau$. In both cases $W(E)$ monotonically increases with energy and reaches zero at $E = E_{sph}$.

At low energies, $E \ll E_{sph}$, the periodic instanton can be approximated by an infinite chain of alternating instantons and anti-instantons [2, 1]. In models which have localized instanton solutions (two-dimensional Abelian Higgs model, quantum mechanics in double well potential, two-dimensional $-\lambda\phi^4$ model) the limit $E \rightarrow 0$ corresponds to $\tau \rightarrow \infty$. Alternatively, in models with no localized instanton (four-dimensional massive $-\lambda\phi^4$ model, $O(3)$ σ -model with the mass term added, Electroweak Theory), $\tau \rightarrow 0$ in the low energy limit². In both cases the dilute gas approximation works and the action of the periodic instanton tends to twice the instanton action, S_0 (from below and from above, respectively, as follows from eq. (3) at $E > 0$).

These two types of behavior at $E \ll E_{sph}$ can be smoothly continued into the region $E \sim E_{sph}$ as shown in Fig.1. In the case I the period τ is always greater than τ_{sph} ; the action monotonically increases from $S_{sph} \equiv E_{sph}\tau_{sph}$ to S_0 as the period changes from τ_{sph} to ∞ (energy changes from E_{sph} to zero). The curve $S(\tau)$ is convex

²In these models the low energy periodic instanton is built out of constrained instantons whose size ρ is stabilized by the interaction with their neighbors in the chain. At $E \rightarrow 0$ both ρ and τ go to zero in such a way that $\rho/\tau \rightarrow 0$ [1].

upward,

$$\frac{\partial^2 S}{\partial \tau^2} = \left(\frac{\partial^2 W}{\partial E^2} \right)^{-1} = \frac{\partial E}{\partial \tau} < 0.$$

Correspondingly, the saddle point (5) is a *maximum* of the exponent in eq. (4) and saturates the thermal rate at temperatures $T = 1/\beta < 1/\tau_{sph}$, so that one has

$$\begin{aligned} \Gamma(\beta) &\sim \exp(-S(\beta)) & \text{at } T < 1/\tau_{sph}, \\ \Gamma(\beta) &\sim \exp(-\beta E_{sph}) & \text{at } T > 1/\tau_{sph}. \end{aligned}$$

Note that this case requires $S_0 > S_{sph}$.

In the case II the period is always less than τ_{sph} . The action monotonically decreases when the period changes from τ_{sph} to zero (energy changes from E_{sph} to zero). The curve $S(\tau)$ is convex downward,

$$\frac{\partial^2 S}{\partial \tau^2} = \left(\frac{\partial^2 W}{\partial E^2} \right)^{-1} = \frac{\partial E}{\partial \tau} > 0.$$

In this case, the saddle point (5) is a *minimum*; the integral in eq. (4) is saturated either by $E = 0$ or by $E = E_{sph}$, depending on the temperature:

$$\begin{aligned} \Gamma &\sim \exp(-S_0) & \text{at } T < E_{sph}/S_0, \\ \Gamma &\sim \exp(-\beta E_{sph}) & \text{at } T > E_{sph}/S_0. \end{aligned}$$

This case requires $S_0 < S_{sph}$.

Do these two cases comprise all possible types of behaviour? The answer is *no*. It is easy to find models where neither case I nor case II can realize. In fact, the Electroweak Theory is one of the examples. It has no localized instantons and shows low energy behaviour of the periodic instanton as in the case II [1]. However, as was first noted in ref.[5], the numerical analysis of the electroweak sphaleron [7] implies that at $M_H > 4M_W$ one has $S_0 > S_{sph}$, which is not consistent with the case II³. This situation is rather common and is due to the fact that low and high energy limits of periodic instanton are determined by different parts of the potential and hence are not tied to each other. Below we present two examples of models where the periodic instanton can be found explicitly and shows the behaviour more complicated than in the cases I and II of Fig.1.

A quantum mechanical model. Consider one-dimensional quantum mechanical model with the Euclidean action

$$S_E = \frac{1}{\lambda} \int d\tau \left(\frac{1}{2} \dot{\phi}^2 + V(\phi) \right).$$

³This anomaly has nothing to do with deformed sphaleron which appears only at $M_H \simeq 12M_W$

The potential $V(\phi)$ is chosen in the form

$$V(\phi) = \frac{4 + \alpha}{12} - \frac{1}{2}\phi^2 - \frac{\alpha}{4}\phi^4 + \frac{\alpha + 1}{6}\phi^6.$$

When $\alpha > -1$ it is of the double well type, i.e. it has two minima at $\phi = \pm 1$ separated by the barrier of the height

$$E_{sph} = \frac{4 + \alpha}{12\lambda}.$$

The main difference with the standard (quartic) case is that the walls of the wells are steeper.

This model possesses exact zero-energy instanton describing quantum tunneling between two wells, in full analogy with conventional double well model. Therefore, at $E \rightarrow 0$ the period of the periodic instanton goes to infinity (this is easy to check by direct calculation, see eq. (6) below). On the other hand, in the vicinity of the sphaleron one finds

$$S(\tau) = E_{sph}\tau + \frac{4\pi\alpha}{3\lambda} \left(1 - \frac{\tau}{2\pi}\right)^2,$$

so that at $\alpha > 0$ the periodic instanton has period *smaller* than $\tau_{sph} = 2\pi$ and the value of action *larger* than the corresponding sphaleron value.

In order to see how the two limiting cases match together one has to find the periodic instanton at intermediate energies. In quantum-mechanical model this can be done in quadratures,

$$\begin{aligned} S(\tau) &= E\tau + \frac{1}{\lambda} \oint dx \sqrt{2(V(x) - \lambda E)} \\ \tau &= \oint \frac{dx}{\sqrt{2(V(x) - \lambda E)}} \end{aligned}$$

The function $S(\tau)$ for $\alpha = 1.5$ is shown in Fig. 2. The periodic instanton indeed exists in some range $\tau < \tau_{sph}$, but at a critical value of period $\tau_c < \tau_{sph}$ the curve $S(\tau)$ has a *wedge* and turns back to large periods. The branches of the wedge are tangent at the bifurcation point, as follows from energy continuity and eq. (3). The curve is convex upward on the lower branch and downward on the upper one.

σ -model. A field theory model with similar behaviour of the periodic instanton can be constructed from the massive $O(3)$ σ -model of ref.[8] by adding the Skyrme term. The model is defined by the Euclidean action

$$S_E = \frac{1}{g^2} \int d^2x \left\{ \frac{1}{2} n_\mu^a n_\mu^a + (1 + n_3) + \frac{\lambda}{8} n_{\mu\nu}^{ab} n_{\mu\nu}^{ab} \right\}, \quad (6)$$

where the n^a is a unit vector in 3-dimensional space $a = 1, 2, 3$, $n_\mu^a = \partial_\mu n^a$ and $n_{\mu\nu}^{ab} = n_{[\mu}^a n_{\nu]}^b$. Without the Skyrme term (i.e. at $\lambda = 0$), the massive σ -model does not




Figure 2: The periodic instanton action as a function of its period in the steep double well model.

have localized instantons and falls into the class II [5]. The approximate solutions, constrained instantons, are unstable and shrink to zero size. At $\lambda \neq 0$, the size of the constrained instanton stabilizes at the value of order $(\lambda \log \lambda)^{1/4}$ and the model (6) acquires a localized instanton solution. It can be found numerically [9]. The double instanton action at $\lambda \ll 1$ is

$$S_0 = \frac{8\pi}{g^2} + O\left(\frac{\lambda^{1/2}}{g^2}\right).$$

Thus, at $\lambda \neq 0$ one expects the low energy behaviour as in the case I.

The Skyrme term does not affect static configurations, so the sphaleron solution coincides with that of $\lambda = 0$ model, where its analytical form is [8]

$$n^a = \left(-2 \frac{\text{sh}(x)}{\text{ch}^2(x)}, 0, -1 + \frac{2}{\text{ch}^2(x)} \right).$$

The energy of the sphaleron equals

$$E_{sph} = \frac{8}{g^2}.$$

The Skyrme term distorts the matrix of the second derivatives of the action, so the negative sphaleron eigenmode and, hence, the critical period depend on λ . It still can be found analytically,

$$u_-^a = \delta_2^a \frac{1}{\text{ch}^\alpha(x)}, \quad (7)$$




Figure 3: (a) The action of the periodic instanton ($\diamond$) in the σ -model with the Skyrme term, as a function of its period. Straight line represents the sphaleron. The behaviour of $S(\beta) = -\ln(\Gamma(\beta))$ is shown by fat line. (b) The energy of the same periodic instanton as a function of the period.

$$\omega_- = (\alpha^2 - 1)^{1/2},$$

where α is the positive root of the equation

$$(1 + 4\lambda)\alpha^2 + \alpha - 4\lambda - 6 = 0.$$

The action of the sphaleron at the critical period is

$$S_{sph} = E_{sph}\tau_{sph} = \frac{16\pi}{\sqrt{3}g^2} + O\left(\frac{\lambda}{g^2}\right).$$

Note that $S_0 < S_{sph}$, which is not compatible with the case I.

The periodic instanton can be found numerically. The results of the calculation at $\lambda = 0.001$ are presented in Fig. 3. As in the case of the steep double well model, there is a wedge at $\tau < \tau_{sph}$.

In both examples, the function $S(\tau)$ is multi-valued, and the period of the periodic instanton is not a monotonic function of energy. This does not affect qualitatively the probability of tunneling at fixed energy. In both models $W(E)$ is a monotonically increasing function of energy; the wedge in the action shows up as zero of second derivative of $W(E)$. However, the existence of the wedge has non-trivial consequences for the thermal rate $\Gamma(\beta)$. The lower branch of the periodic instanton dominates $\Gamma(\beta)$ at temperatures $T < T_* = 1/\tau_*$, where τ_* is determined by the intersection of the

lower branch with the line $E_{sph}\tau$. Note that $T_* > T_{sph}$. At higher temperatures, the rate is controlled by the sphaleron,

$$\begin{aligned}\Gamma(\beta) &\sim \exp(-S(\beta)) && \text{at } T < 1/\tau_*, \\ \Gamma(\beta) &\sim \exp(-\beta E_{sph}) && \text{at } T > 1/\tau_*.\end{aligned}$$

Thus, in a sense, the two examples considered above are intermediate between cases I and II.

General properties of periodic instanton bifurcations. The wedge in the action that was found in the above examples is likely to be rather common phenomenon. Moreover, it may be considered as a particular kind of bifurcations of the periodic instanton solution under the change of parameters (period τ here) and should be treated on equal footing with the bifurcation point at $\tau = \tau_{sph}$ where the periodic instanton merges to the sphaleron. Since the behaviour of solutions near bifurcation points is governed by global properties of the action, its general features can be analyzed without referring to particular model.

Consider the hessian of the action for a given periodic instanton I . In general, the periodic instanton is not a local minimum of the action, so the hessian has a finite number $\mu(I)$ of negative eigenvalues. Generally, it has also zero modes, corresponding to global symmetries of the model, but we consider them pre-eliminated, so that for generic values of parameters the hessian is supposed to be non-degenerate and, hence, $\mu(I)$ is the Morse index of the solution I . The determinant of the hessian gives pre-exponential factor in eq. (2), which is ignored throughout the paper. However, it contains an important piece of information about the leading approximation. Namely, each negative mode of the hessian contributes into eq. (2) a factor i . The path integral representation for $P(E)$ contains also integrations over discrete variables which finally reduce to integration over the period τ [1] (so that eq. (3) is in fact the saddle point equation). This integration may give another factor i . Thus, the requirement that $P(E)$ is real translates into equation

$$\left[\text{sign} \frac{\partial E}{\partial \tau} \right] (-1)^{\mu(I)} = 1, \quad (8)$$

where we made use of the relation $\partial^2 S / \partial^2 \tau = \partial E / \partial \tau$. Eq.(8) should be treated as a selection rule for solutions which can contribute into $P(E)$. According to this rule, the periodic instantons of types I and II should have an odd and even number of negative modes, respectively. As is shown below, this is indeed the case. Similar statement applies to the lower and upper branches of the wedge.

When several solutions merge together one can define the degree d of the bifurcation point as sum of $(-1)^{\mu(I)}$ over all merging solutions:

$$d = \sum_I (-1)^{\mu(I)}.$$

This number is a topological invariant, which has the same value before and after bifurcation. Actually, the low dimensional geometric intuition implies that the number of negative modes on merging branches monotonically increases with their action.

As an example consider the wedge discussed above. Before the bifurcation point ($\tau < \tau_c$) there are no solutions, so $d = 0$. Therefore, at $\tau > \tau_c$ the two emerging solutions must obey $\mu(I_1) + \mu(I_2) = \text{odd}$ (and, therefore, should have opposite signs of $\partial E / \partial \tau$), i.e. the lower and upper branches should have an odd and even number of negative modes, respectively. This fact can be easily checked in quantum mechanics by direct numerical diagonalization of the discretized hessian; for field models this statement is less trivial.

Another application of the conservation of d is the bifurcation of the sphaleron at $\tau = \tau_{sph}$. The sphaleron solution has exactly one static negative mode. Consider time dependent modes of the hessian

$$H = -\partial_\tau^2 + H_{stat}$$

in the vicinity of the sphaleron. Here H_{stat} is the time-independent part of the hessian. The static eigenmodes of the sphaleron diagonalize H_{stat} ,

$$H_{stat} = \sum \omega_n^2 u_n \otimes u_n.$$

Therefore, the full hessian has eigenmodes

$$U_{mn}(t, x) = \cos \frac{2\pi m t}{\tau} u_n(x)$$

and eigenvalues

$$\Lambda_{mn} = \omega_n^2 + \left(\frac{2\pi m}{\tau} \right)^2.$$

The static negative mode of the sphaleron corresponds to $n = 0$, so that $\omega_0^2 = -\omega_-^2 < 0$. We see that at $\tau < \tau_{sph} = 2\pi/\omega_-$ the full hessian has exactly one negative mode $\Lambda_{00} = -\omega_-^2$, but at $\tau > \tau_{sph}$ another negative mode appears with the eigenvalue

$$\Lambda_{10} = -\omega_-^2 + \left(\frac{2\pi}{\tau} \right)^2 < 0.$$

In order to satisfy the conservation of d at $\tau = \tau_{sph}$ we have to assume the existence of at least *two* periodic instanton solutions, both merging or splitting when τ passes through τ_{sph} . Moreover, in the case of splitting these solutions should have one negative mode each, while in the case of merging two negative modes each. We indeed have exactly two periodic instantons differing by translation in time by $\tau/2$, so this analysis not only proves the existence of periodic instanton in the vicinity of the sphaleron (this fact is non-trivial in models with multiple degrees of freedom), but also predicts the number of their negative modes.




Figure 4: (a) Expected behaviour of the periodic instanton action, $S(\tau)$, in the Electroweak theory at $4M_W < M_H < 12M_W$. Straight line represents the sphaleron. (b) Corresponding function $S(\beta)$ (fat line) which determines the thermal rate $\Gamma \sim \exp[-S(\beta)]$.

To summarize this discussion, the behaviour of periodic instantons is governed by a set of simple rules: i) The action always increases with the period, as follows from eq. (3). ii) The energy monotonically decreases as moving along the curve of periodic instantons from the sphaleron. iii) The curve is convex upward or downward at $\partial E/\partial\tau$ negative or positive, respectively. iv) The number of negative modes is correlated with the sign of $\partial E/\partial\tau$ so as to obey eq. (8). v) At bifurcation points the degree d must conserve and the solution with larger action should have more negative modes.

In conclusion, let us apply these rules to the periodic instanton in the Electroweak Theory at $12M_W > M_H > 4M_W$. In this case one has $S_0 > S_{sph}$, while the low energy behaviour of the periodic instanton follows the case II. The simplest way to reconcile these two facts is shown in Fig. 4a (if the behaviour of the periodic instanton near the sphaleron is as in the case II, it must have at least two wedges). The lower branch (consisting of two identical periodic instantons which are shifted by $\tau/2$) has one negative mode, while the upper branch has two negative modes. This has an implication in the calculation of the thermal rate: at temperatures T satisfying $\tau_{sph} < 1/T < \tau_*$ the latter is saturated by the lower branch of periodic instanton, see Fig. 4b. This picture survives up to $M_H \sim 12M_W$, where the sphaleron itself bifurcates [7].

Acknowledgements. The authors are grateful to V. A. Rubakov for numerous discussions at different stages of this work. The work is supported in part by Award No. RP1-187 of the U.S. Civilian Research & Development Foundation for the Independent States of the Former Soviet Union (CRDF), and by Russian Foundation for

Basic Research, grant 96-02-17804a.

References

- [1] S. Yu. Khlebnikov, V. A. Rubakov and P. G. Tinyakov, *Nucl. Phys.* **B367** (1991) 334.
- [2] A.D. Linde, *Phys. Lett.* **100B** (1981) 37, *Nucl. Phys.* **B216** (1983) 421.
- [3] N. S. Manton, *Phys. Rev.* **D28** (1983) 2019; F. R. Klinkhamer and N. S. Manton, *ibid.* **30** (1984) 2212.
- [4] V. V. Matveev, *Phys. Lett.* **B304** (1993) 291.
- [5] S. Habib, E. Mottola and P. G. Tinyakov, *Phys. Rev.* **D54** (1996) 7774.
- [6] A. N. Kuznetsov and P. G. Tinyakov, *Mod. Phys. Lett.* **A11** (1996) 479; “*False Vacuum Decay Induced by Particle Collisions*”, submitted to *Phys. Rev. D*, hep-ph/9703265.
- [7] L. G. Yaffe, *Phys. Rev.* **D40** (1989) 3463.
- [8] E. Mottola and A. Wipf, *Phys. Rev.* **D39** (1989) 588.
- [9] B. M. A. G. Piette, H. J. W. Müller-Kirsten, D. H. Tchrakian and W. J. Zakrewski, *Phys. Lett.* **B320** (1994) 294.